

bonjela teething gel

Oromucosal Gel
Choline salicylate 8.714%
Cetalkonium chloride 0.010%

PACKAGE LEAFLET: INFORMATION FOR THE USER

Read all of this leaflet carefully because it contains important information for you.

This medicine is available without prescription. However, you still need to use Bonjela Teething Gel carefully to get the best results from it. Keep this leaflet. You may need to read it again. Ask your pharmacist if you need more information or advice. You must contact a doctor if your symptoms worsen or do not improve after 7 days.

If any of the side effects gets serious, or if you notice any side effect not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Bonjela Teething Gel is and what it is used for
2. Before you use Bonjela Teething Gel
3. How to use Bonjela Teething Gel
4. Possible side effects
5. How to store Bonjela Teething Gel
6. Further information

1. WHAT BONJELA TEETHING GEL IS AND WHAT IT IS USED FOR

Bonjela Teething Gel is a clear gel, which is applied to the inside of the mouth or gums. It contains choline salicylate (a pain killer and anti-inflammatory) which soothes the sore areas in the mouth, and antiseptic called cetalkonium chloride, which kills bacteria and helps prevent infection.

Bonjela Teething Gel can be used to relieve pain and reduce inflammation and help fight infection in infant teething, mouth ulcers, cold sores, and sore spots due to dentures in adults and orthodontic devices in children.

2. BEFORE YOU USE BONJELA TEETHING GEL

Do not use Bonjela Teething Gel:

- if you are allergic (hypersensitive) to the active ingredients or any of the ingredients listed in section 6.
- if you are currently taking aspirin or other salicylate, except under direction of a doctor
- in children under 4 months
- If you have an active or history of recurrent stomach ulcer
- if you are currently suffering from chicken pox or flu particularly in children and adolescents under 16 years of age

Take special care with Bonjela Teething Gel

- if you are pregnant or breastfeeding, please consult your doctor before use
- if there is no improvement after 7 days, consult your doctor
- especially in children, as frequent application may give rise to salicylate poisoning.

Taking other medicines

Salicylates may enhance the effect of blood thinning medicines (anticoagulants) and can decrease the effect of medicines to treat gout (uricosurics) such as probenecid.

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Pregnancy and breast-feeding

If you are pregnant, planning to become pregnant, or breastfeeding, speak to your doctor or your pharmacist.

Important information about some of the ingredients of Bonjela Teething Gel

Bonjela Teething Gel contains a small amount of ethanol (alcohol), less than 100mg per dose.

Contains glycerol, may be harmful in high doses and can cause headache, stomach upset and diarrhoea.



3082008

Pharmacode Reads This Way

First Bar

Last Bar

3. HOW TO USE BONJELA TEETHING GEL

Bonjela Teething Gel is for oral topical use on the mouth and gums. Check that the cap seal is unbroken before first using this product. Apply with a clean finger.

Children (from 4 months) – massage a pea sized amount of gel onto the sore area, not more than once every 3 hours. Do not apply more than six times in 24 hours.


Although this product is for babies, it may also be used in Adults the following administration applies;

Adults – Massage approximately 1cm of the gel onto the sore area, not more than once every 3 hours.

Orthodontic Devices – Apply to sore area as described above, not more than once every 3 hours.

Dentures Irritation – Apply to the gums and leave at least 30 minutes before re-insertion of the dentures. Do not apply directly to the dentures.

DO NOT EXCEED THE STATED DOSE



CHILDREN OVER 4MTHS

ADULTS

If your symptoms do not improve after 7 days of treatment you should contact your doctor.

If you use more Bonjela Teething Gel than you should

Contact your doctor or pharmacist. Symptoms of overdose include nausea, vomiting, fever and disorientation. It is especially important in infants to follow the dosage instructions carefully.

If you forget to use Bonjela Teething Gel

Apply a new dose as described above. Do not take a double dose to make up for a forgotten dose. If you have any further questions on the use of this product, ask your doctor or pharmacist.

Ink &
Varnish Free

4. POSSIBLE SIDE EFFECTS

Like all medicines, Bonjela Teething Gel can cause side effects, although not everybody gets them. Allergic reactions to salicylates such as an asthma attack or breathing difficulties may occur. Other reactions such as inflammation, burns, numbing or stinging of the mouth or gums may occur. There have been post-marketing reports of cases of suspected Reyes syndrome and possible salicylate toxicity in children who had a history of use of choline salicylate gels but causality has not been established.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor, pharmacist or nurse: You can report side effects directly via HPR Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie

5. HOW TO STORE BONJELA TEETHING GEL

Do not store above 25°C. Keep out of the reach and sight of children.

Do not use Bonjela Teething Gel after the expiry date which is stated on the carton and tube. The expiry date refers to the last day of that month. Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

What Bonjela Teething Gel contains

The active ingredients are choline salicylate 8.714% w/w and cetalkonium chloride 0.01% w/w. The other ingredients are ethanol, glycerol, levomenthol, hypermellose 4500, star anise oil, saccharin sodium and purified water.

What Bonjela Teething Gel looks like and contents of the pack

Each pack contains a 15g tube of clear, colourless, aniseed flavoured, oromucosal gel.

Marketing Authorisation Holder and Manufacturers

PA Holder: Reckitt Benckiser Ireland Ltd., 7 Riverwalk, Citywest Business Campus, Dublin 24, Ireland. Manufacturers:

Reckitt Benckiser (UK) Ltd., Dansom Lane, Hull, HU8 7DS, UK and RB NL Brands B.V. Schiphol Blvd 207, 1118 BH Schiphol, NL.

For product queries please call: 01-6305429

Date of revision of text: December 2018.



3082008