


NUROFEN®

5% W/W Gel

Ibuprofen



Read all of this leaflet carefully because it contains important information for you.

This medicine is available without prescription.

However, you still need to use Nurofen 5% w/w Gel carefully to get the best results from it.

Keep this leaflet. You may need to read it again.

Ask your pharmacist if you need more information or advice.

You must contact a doctor if your symptoms worsen or do not improve after 7 days.

If any of the side-effects gets serious, or if you notice any side-effect not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Nurofen 5% w/w Gel is and what it is used for
2. Before you use Nurofen 5% w/w Gel
3. How to use Nurofen 5% w/w Gel
4. Possible side-effects
5. How to store Nurofen 5% w/w Gel
6. Further Information

1. WHAT NUROFEN 5% W/W GEL IS AND WHAT IT IS USED FOR?

Nurofen 5% w/w Gel contains the active ingredient ibuprofen which belongs to a group of medicines known as non steroidal anti-inflammatory drugs (NSAIDs).

It is gently rubbed into the skin

- to relieve pain, reduce local heat, swelling and inflammation
- in conditions such as backache, rheumatic and muscular pain, sprains, strains and sports injuries.

2. BEFORE YOU USE NUROFEN 5% W/W GEL:

Do not use Nurofen 5% w/w Gel if you:

- are **allergic** to ibuprofen or any of the ingredients of Nurofen 5% w/w Gel listed in Section 6.
- are **allergic** to aspirin or other pain relievers known as non steroidal anti-inflammatory drugs (NSAIDs)
- have ever had an **asthma attack, rash or runny nose after taking aspirin** or any NSAIDs

- have broken or damaged skin
- are **under 12 years of age**
- are in the third trimester of pregnancy

Consult your doctor or pharmacist before taking Nurofen 5% w/w Gel if you

- have asthma or have suffered from asthma
- are **taking aspirin, ibuprofen or other NSAID** pain relievers
- Are pregnant

Warnings and Precautions

Nurofen Gel is applied to the skin, there is less risk of the complications that sometimes occur when ibuprofen (or other NSAIDs) is taken by mouth. However, in rare cases you may be at risk:

- if you have a stomach ulcer (also called peptic or gastric ulcer)
- if you have kidney problems
- if you have liver problems
- if you have ever had asthma
- if you have ever had intestinal inflammation or hemorrhage


Take special care with Nurofen 5% Gel

- Use it only on the skin
- Keep the gel away from the eyes, nose and mouth
- Do not use it under waterproof dressings
- If you develop a rash after using the gel stop using it any further
- Protect treated areas from direct artificial or natural light to avoid any sensitivity reaction, e.g. a rash


Taking other medicines

Some medicines that are anti-coagulants (i.e. thin blood/prevent clotting, e.g. aspirin/ acetylsalicylic acid, warfarin, ticlopidine), some medicines that reduce high blood pressure (ACE-inhibitors such as captopril, beta-blockers such as atenolol or angiotensin II receptor antagonists such as losartan)

3060584



Pharmacode Height = 8mm
Pharmacode Width = 8mm minimum
6mm clear area left and right
6mm clear area above and below
Reverse of Pharmacode is print free



Pharmacode Height = 8mm
Pharmacode Width = 8mm minimum
6mm clear area left and right
6mm clear area above and below
Reverse of Pharmacode is print free

1st Fold

4th Fold

3rd Fold

2nd Fold

and other medicines may affect or be affected by treatment with ibuprofen. You should therefore always seek the advice of a healthcare professional before you use ibuprofen with other medicines.

Pregnancy and breast-feeding

You should not use Nurofen 5% Gel if you are pregnant or breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

3. HOW TO USE NUROFEN 5% W/W GEL:

For cutaneous use (use on the skin) only. When first opening the tube, check that the tube seal is not broken. To break seal, press against the point hidden in the top surface of the cap.

Adults and children 12 years and over

Squeeze 50 to 125mg (4 to 10cm) of gel from the tube to the affected area and massage gently until absorbed.

Leave four hours between doses and do not apply more than 4 times in any 24 hour period.

Hands should be washed immediately after use. Do not exceed the stated dose.

Children under 12 years: Do not use on children under 12 years of age.

Do not

- Apply it more than 4 times in 24 hours.
- Use it on broken or inflamed skin, on the lips, genital area or near the eyes.
- Cover the area with bandages, dressings or plasters.

This gel is for **external use only**.

Length of treatment

If your symptoms worsen or if there is no improvement after 7 days treatment, talk to your doctor.

Stop using the gel if you develop **a rash** (see section 4).

If the gel is swallowed by mistake

If the gel is swallowed, rinse the mouth out with water. Swallowing the gel may cause an upset stomach. If you are concerned, talk to your doctor or pharmacist. If a child swallows the gel, talk to your doctor straight away.

4. POSSIBLE SIDE EFFECTS:

Like all medicines, Nurofen 5% w/w Gel can cause side-effects, although not everyone gets them.

Seek medical help at once if you develop any of the following rare reactions.

- unexplained wheeziness, shortness of breath, onset of asthma, swelling of the face, lips or tongue, difficulty swallowing or breathing
- serious skin reactions such as itchy rashes, inflammation, swelling, peeling or cracking of the skin, raised red patches with blistering.
- kidney problems such as swelling or decreased urine out put

Tell your doctor if you are concerned about any of the following effects or have any other effects.

- mild reddening, tingling or itching where you have applied the gel
- stomach pain or indigestion.
- sudden constriction of the airways leading to difficulties breathing (particularly in people who have a history of asthma or allergic problems).

5. HOW TO STORE NUROFEN 5% W/W GEL?

KEEP OUT OF REACH AND SIGHT OF CHILDREN
Do not use after the expiry date printed on the carton and the tube.

Do not store above 25°C.

6. FURTHER INFORMATION

What Nurofen 5% w/w Gel contains

The active substance in Nurofen 5% w/w Gel is ibuprofen 5% w/w. The other ingredients are purified water, isopropyl alcohol, hydroxyethylcellulose, benzyl alcohol, sodium hydroxide (E524).

What Nurofen 5% w/w Gel looks like and contents of the pack

Nurofen 5% w/w Gel is a clear, colourless, smooth gel. It is available in a 30g tube.

Marketing Authorization Holder and Manufacturers

PA Holder: Reckitt Benckiser Ireland Ltd, 7 Riverwalk, Citywest Business Campus, Dublin 24, Ireland.
Manufacturers: Farmasierra Manufacturing S.L., Ctra de Irún, Km 26,200, 28700 San Sebastian de los Reyes, Madrid, Spain.

Reckitt Benckiser Healthcare UK Limited, Dansom Lane, Hull, HU87DS, UK and RB NL Brands B.V. Schiphol Blvd 207, 1118 BH Schiphol, NL

For product enquiries, please telephone (01) 6305429. This leaflet was updated in February 2020

3060584