

IN THIS LEAFLET

SUGAR FREE

680mg/80mg CHEWABLE TABLETS

Calcium carbonate

Magnesium carbonate

Read all of this leaflet carefully because it contains

important information for you.

This medicine is available without prescription. However,

you still need to take Rennie® Sugar Free tablets carefully

to get the best results from them. Keep this leaflet, you

may need to read it again. Ask your pharmacist if you need more information or advice. You must contact a

doctor if your symptoms worsen or do not improve. If you

have any unusual effects after taking Rennie Sugar Free,

tell your doctor or pharmacist.

1. What is Rennie® Sugar Free and what is it used for?

2. Before you take Rennie® Sugar Free

3. How to take Rennie® Sugar Free 4. Possible side effects

5. How to store Rennie® Sugar Free 6. Further information 1. WHAT IS RENNIE® SUGAR FREE AND WHAT IS IT USED

FOR? Rennie® Sugar Free is a mint flavour antacid tablet which relieves stomach upset due to excess acid and heartburn (acid indigestion).

2. BEFORE YOU TAKE RENNIE® SUGAR FREE Do not take Rennie® Sugar Free:

 If you have severe kidney disease or suffer from kidney stones.

If you have high calcium levels in the blood, kidneys or urine.

Take special care with Rennie® Sugar Free: As with other antacid products, taking these tablets can

mask the symptoms of other, more serious, underlying medical conditions. Do not exceed the stated dose.

If symptoms persist after seven days consult your doctor. Patients aged 55 years or older with unexplained and

persistent heartburn should contact their doctor before

taking Rennie® Sugar Free. Rennie® is designed to help relieve pain and discomfort associated with excess stomach acid. Pain/discomfort

associated with cardiac conditions may on occasion mimic

indigestion; such pain will not respond to Rennie®, therefore it is recommended that medical advice be sought if the pain does not go away. -

Consult your doctor before taking Rennie® Sugar Free if you have any kidney problems. This product contains sorbitol (E420). If you have been told by your doctor that you have an intolerance to some sugars, contact your

Taking other medicines: Please tell your doctor or pharmacist if you are taking. or have recently taken any other medicines, including medicines obtained without a prescription. · If you are taking antibiotics (tetracyclines, quinolones),

doctor before taking this medicinal product.

medication for heart failure (e.g digoxin), thiazide diuretics (water tablets), levothyroxine (a thyroid medication) or other prescribed medicines (e.g. fluorides, phosphates, iron containing products, eltrombopag)

speak to your doctor or pharmacist before taking Rennie® Sugar Free because it can affect how these medicines work. · To maximise the benefit of all medicines being used,

If you have low phosphate levels in the blood. take Rennie® Sugar Free 1 to 2 hours after taking any If you are allergic (hypersensitive) to any of the other medicines. ingredients (listed in Section 6. Further information).

Adults: 1 or 2 tablets to be sucked or chewed as required, If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed up to a maximum of 11 tablets a day. in this leaflet. You can also report side effects directly via Children and adolescents below age 18: Not HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2:

3. HOW TO TAKE RENNIE® SUGAR FREE

recommended.

Pregnancy and breast feeding: Ask-your doctor or

pharmacist for advice before taking any medicine. Rennie® Sugar Free can be used during pregnancy if taken as instructed and if prolonged use of high doses is avoided. Rennie® Sugar

Free can be used while breast-feeding if taken as instructed. As Rennie® Sugar Free provides a substantial amount of calcium in addition to dietary calcium intake pregnant women should strictly limit their use of Rennie® Sugar Free

tablets to the maximum recommended daily dose and avoid concomitant, excessive intake of milk (1 litre contains up to 1.2g elemental calcium) and dairy products.

If you take more Rennie® Sugar Free than you should: Drink plenty of water and consult your doctor or pharmacist. Symptoms of an overdose include nausea and vomiting, constipation, tiredness, increased urine production, increased thirst, dehydration and abnormal muscular weakness. If you have any questions on the use of this product, ask

your doctor or pharmacist. 4. POSSIBLE SIDE EFFECTS Like all medicines, Rennie® Sugar Free can cause side effects, although not everybody gets them. Side effects are unlikely at the recommended dose.

Very rarely severe allergic reactions e.g. rashes, swelling or shortness of breath may occur. If you experience these reactions stop treatment and contact a doctor or emergency department immediately.

Long term use of high doses can cause high calcium and magnesium levels in the blood, nausea, vomiting, stomach discomfort, diarrhoea and milk-alkali syndrome, especially in people with kidney conditions. This can cause nausea and vomiting, constipation, headache, alterations

in taste sensations and muscular weakness.

Marketing Authorisation Holder and Manufacturer: Marketing Authorisation Holder: Bayer Limited, The Atrium, Blackthorn Road, Dublin 18.

of the pack:

Reporting of side effects

Tel: +353 1 6764971; Fax: +353 1 6762517.

information on the safety of this medicine.

5. HOW TO STORE RENNIE® SUGAR FREE

above 25°C and keep in the original pack.

to the last day of that month.

protect the environment.

6. FURTHER INFORMATION

What Rennie® Sugar Free contains:

Each tablet contains the active ingredients: calcium

The other ingredients are sorbitol (E420), magnesium

spearmint flavour, liquid paraffin, saccharin sodium.

Rennie® Sugar Free are white square tablets

engraved "RENNIE" on both faces.

carbonate 680mg and heavy magnesium carbonate 80mg.

stearate, potato starch, pregelatinised maize starch, talc,

What Rennie® Sugar Free looks like and the contents

Rennie® Sugar Free are available in packs of 24, 48, and 96

chewable tablets. Not all pack sizes may be marketed

Website: www.hpra.ie; E-mail: medsafety@hpra.ie.

By reporting side effects, you can help provide more

Keep out of the reach and sight of children. Do not store

Do not use Rennie® Sugar Free after the expiry date which

is stated on the carton and blister. The expiry date refers

Medicines should not be disposed of via waste water or

household waste. Ask your pharmacist how to dispose of

medicines no longer required. These measures will help to

Label last revised: February 2015. Bayer 101748 / 0315IE