

[illegible]

Do not exceed the stated dose. If you are having digestive surgery, you should mouthwash before and after surgery as directed by your dentist. You should use the mouthwash for as long as directed. Do not swallow the mouthwash. See below. For the treatment of red, swollen or inflamed oral tissues, you should use the mouthwash for about one month. You will probably also need to use the mouthwash for one month. For mouth ulcers, denture stomatitis, and thrush, continue using the mouthwash for two days after the symptoms have disappeared, up to a maximum of 14 days. If you are using the mouthwash for denture stomatitis, you should also thoroughly clean your dentures. The recommended dose of the mouthwash for fifteen minutes twice daily.

If your symptoms persist after you have used the mouthwash for the recommended time,

using the mouthwash without any further problems.

If you experience or irritation of the mouth, stop using the mouthwash and tell your dentist, doctor or pharmacist.

Very occasionally you may get some swelling or a rash in the salivary glands when using the mouthwash. This should disappear once you have stopped using it.

Very rarely severe allergic reactions may occur (anaphylaxis). If you get a rash, swelling of the face, tongue, throat, face or have difficulty breathing, you should see a doctor immediately.

Some people may experience some of the following symptoms, or you experience any other unusual symptoms, or you experience any of the following symptoms, or doctor or pharmacist at once.

Reporting of side effects: If you get any side effects, talk to your doctor, pharmacist or nurse.

Any such information on the product or its use, directly via HPA, Pharmacovigilance, medwatch@hpa.gov.uk, Tel: +33 16749741 Fax: +333 1 6762571, Web: medwatch.hpa.gov.uk

For more information on the safety of this medicine, see **5. How to store Gaseodyn® Anisised**

Do not use the mouthwash after the expiry date (EXP) shown on the pack. The expiry date is the date on the mouthwash container above 25° C. Store in the original container and keep out of the reach of children.